

Clinical profile and predictors of in-hospital mortality among older patients admitted for COVID-19

Running head: COVID-19 in older patients.

Víctor Manuel Becerra-Muñoz^{1*}, Iván J. Núñez-Gil^{2*}, Charbel Maroun Eid³,
Marcos García Aguado⁴, Rodolfo Romero⁵, Jia Huang⁶, Alba Mulet⁷, Fabrizio
Ugo⁸, Francesco Rametta⁸, Christoph Liebetrau⁹, Alvaro Aparisi¹⁰,
Inmaculada Fernández-Rozas¹¹, María C. Viana-Llamas¹², Gisela Feltes¹³,
Martino Pepe¹⁴, Luis A. Moreno-Rondón¹⁵, Enrico Cerrato¹⁶, Sergio
Raposeiras-Roubín¹⁷, Emilio Alfonso¹⁸, Ana Carrero-Fernández¹⁹, Luís
Buzón-Martín²⁰, Mohammad Abumayyaleh²¹, Adelina Gonzalez²², Antonio
Fernández Ortiz², Carlos Macaya², Vicente Estrada², Cristina Fernández-
Pérez², Juan José Gómez-Doblas¹.
On behalf of HOPE COVID-19 investigators.

AFFILIATION

- 1) Unidad de Gestión Clínica Área del Corazón, Instituto de Investigación Biomédica de Málaga (IBIMA), Hospital Universitario Virgen de la Victoria, Universidad de Málaga (UMA), Centro de Investigación Biomédica en Red de Enfermedades Cardiovasculares (CIBERCV), Málaga, Spain.
- 2) Hospital Clínico San Carlos. Universidad Complutense de Madrid, Instituto de Investigación Sanitaria del Hospital Clínico San Carlos (IdISSC), Centro de Investigación Biomédica en Red de Enfermedades Cardiovasculares (CIBERCV), Madrid, Spain.
- 3) Hospital Universitario La Paz. Instituto de Investigación Hospital Universitario La Paz (IdiPAZ), Madrid, Spain.
- 4) Hospital Universitario Puerta de Hierro, Majadahonda, Spain.
- 5) Hospital Universitario Getafe, Madrid, Spain.
- 6) The Second Affiliated Hospital of Southern University of Science and Technology, Shenzhen, China.
- 7) Hospital Clínico Universitario, Incliva, Universidad de Valencia, Valencia, Spain
- 8) Sant'Andrea Hospital; Vercelli, Italy
- 9) Kerckhoff Heart and Thorax Center, Bad Nauheim, Germany
- 10) Hospital Clínico Universitario de Valladolid, Valladolid, Spain.
- 11) Hospital Severo Ochoa, Leganés, Spain.
- 12) Hospital Universitario Guadalajara, Guadalajara, Spain.
- 13) Hospital Nuestra Señora de América, Madrid, Spain.
- 14) Azienda ospedaliero-universitaria consorziale policlinico di Bari, Italy.
- 15) Hospital General del norte de Guayaquil IESS Los Ceibos, Guayaquil, Ecuador.
- 16) San Luigi Gonzaga University Hospital, Rivoli, Turin. Italy.
- 17) University Hospital Álvaro Cunqueiro, Vigo. Spain.
- 18) Instituto de Cardiología y Cirugía Cardiovascular, Havana, Cuba.

- 19) Hospital Universitario Príncipe de Asturias, Alcalá de Henares, Spain
- 20) Hospital Universitario de Burgos, Burgos, Spain.
- 21) First Department of Medicine, Medical Faculty Mannheim, University Heidelberg, Mannheim, 68167, Germany, DZHK (German Center for Cardiovascular Research), Partner Site, Heidelberg-Mannheim, Mannheim, Germany.
- 22) Hospital Universitario Infanta Sofía. San Sebastián de los Reyes, Madrid, Spain.

Address for correspondence

Víctor M. Becerra Muñoz, MD, PhD . Department of Cardiology, Hospital Clínico Universitario Virgen de la Victoria, Málaga. Campus de Teatinos, s/n, 29013, Malaga, Spain.
Tel./fax: +34 646790599; E-mail: vmbecerram@gmail.com.

(*) These authors equally contributed to this manuscript.

ABSTRACT

Background. The coronavirus disease 2019 (COVID-19) is characterized by poor outcomes and mortality, particularly in older patients.

Methods. Post-hoc analysis of the international, multicentre, 'real-world' HOPE COVID-19 registry. All patients aged ≥ 65 years hospitalised for COVID-19 were selected. Epidemiological, clinical, analytical and outcome data were obtained. A comparative study between two age subgroups, 65-74 and ≥ 75 years, was performed. The primary endpoint was all cause in-hospital mortality.

Results. 1520 patients aged ≥ 65 years (60.3% male, median age of 76 [IQR 71-83] years) were included. Comorbidities such as hypertension (69.2%), dyslipidemia (48.6%), cardiovascular diseases (any chronic heart disease in 38.4% and cerebrovascular disease in 12.5%), and chronic lung disease (25.3%) were prevalent, and 49.6% were on ACEI/ARBs. Patients aged 75 years and older suffered more in-hospital complications (respiratory failure, heart failure, renal failure, sepsis) and a significantly higher mortality (18.4 vs. 48.2%, $p < 0.001$), but fewer admissions to intensive care units (11.2 vs. 4.8%). In the overall cohort, multivariable analysis demonstrated age ≥ 75 (OR 3.54), chronic kidney disease (OR 3.36), dementia (OR 8,06), peripheral oxygen saturation at admission < 92 % (OR 5.85), severe lymphopenia ($< 500/\text{mm}^3$) (OR 3.36) and qSOFA (Quick Sequential Organ Failure Assessment Score) > 1 (OR 8.31) to be independent predictors of mortality.

Conclusion. Patients aged ≥ 65 years hospitalised for COVID-19 had high rates of in-hospital complications and mortality, especially among patients

75 years or older. Age ≥ 75 years, dementia, peripheral oxygen saturation $< 92\%$, severe lymphopenia and qSOFA scale > 1 were independent predictors of mortality in this population.

KEY WORDS: Coronavirus disease 2019, SARS-CoV-2, older adults, comorbidities.

Key points:

- It is known that COVID-19 is characterized by high morbidity and mortality, particularly in older patients.
- In our study, patients aged 65 years or older had high rates of in-hospital complications and mortality.
- Age ≥ 75 years, dementia, peripheral saturation of O₂ $< 92\%$, severe lymphopenia and qSOFA scale > 1 were predictors of mortality.

INTRODUCTION

On 11th March, 2020 the World Health Organization (WHO) declared the novel coronavirus disease 2019 (COVID-19), a condition caused by the Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2), as a pandemic. [1]

COVID-19 is characterized by poor outcomes and mortality, especially among older patients. [2] In fact, from the beginning of the epidemic, older age has been identified as an important risk factor for disease severity, with increasing rates of mortality across every decade of life. [3-6]

This may be a consequence of poorer previous health status, with higher prevalence of preexisting comorbidities and a higher degree of frailty. [7] Whether worse quality of care in settings where healthcare systems were overwhelmed may also contributed to the observed poor prognosis is unclear. As older patients represent a vulnerable population, comprehensive data are needed in order to improve health care pathways in the context of the COVID-19 pandemic.

Herein, our first objective was to provide a detailed description of clinical characteristics, initial symptoms and management of a cohort of 65 years or older individuals hospitalised for COVID-19. Furthermore, we aimed to present outcomes and to investigate predictors of in-hospital mortality in this highly susceptible population. Finally, we assessed the differences in terms of baseline characteristics, prognosis and treatment among patients 75 years or older.

METHODS

Please see appendix 2.

RESULTS

1520 patients aged ≥ 65 years were included in the study (Appendix 1g). 916 (60.3%) were male and median age was 76 (IQR 71-83) years (Appendix 1h). The most prevalent comorbidities were hypertension (69.2%), dyslipidemia (48.6%), cardiovascular diseases (any heart disease in 38.4% and cerebrovascular disease in 12.5%), and chronic lung disease (25.3%). In the overall cohort, 49.6% were on ACEI/ARBs and 19.3% on oral anticoagulation. Median Charlson Comorbidity Index (CCI) was 1 (IQR 1-3) and the median duration of hospitalization in those patients who survived was 8 (IQR 4-12) days. Clinical baseline characteristics are shown in Table 1 (summary) and Appendix 1a.

At admission, blood tests demonstrated elevated C-reactive protein levels in 92.2% of patients, as well as elevated levels of lactate dehydrogenase (79.2%), D-dimer (72.6%), ferritin (61.2%), transaminases (42.7%) and troponin (21.8%). Severe lymphopenia ($< 500/\text{mm}^3$) was observed in 18.1% of patients.

In-hospital complications and treatment

10.7% of patients had a qSOFA at admission of 2-3 points. During hospitalization, 369 (24.8%) patients developed renal failure (KDIGO stage 1 acute kidney injury or worse), 902 (60.1%) patients developed

respiratory failure, 143 (9.7%) suffered heart failure, 248 (16.7%) suffered sepsis and 342 (23.5%) developed systemic inflammatory response syndrome (SIRS). In addition, 38 patients (2.6%) had an episode of relevant bleeding during hospitalization and 19 (1.3%) an embolic event. Hydroxychloroquine, antibiotics (e.g. Azithromycin) and antiviral drugs were frequently used (82.2, 78.7 and 59.1%, respectively). Almost 80% required oxygen during admission and 6.4% used mechanical ventilation. A detailed description can be found in Table 1.

Patients aged 75 years or older

A comparison between patients aged 65-74 and ≥ 75 years in terms of baseline characteristics, baseline medication, symptoms, in-hospital outcomes and therapies is shown in Table 2 (Summary) and Appendix 1b, S2. Patients aged 75 years or older had more comorbidities such as hypertension, chronic kidney disease, chronic cardiac diseases and dementia, with higher CCI (1.4 vs 2.0 points). Consequently, almost all types of medication were more prevalent at the point of hospital admission. Apart from dyspnea and fatigue, common presenting symptoms of COVID-19 were less prevalent in the older age group' (e.g. fever 82 vs. 74.2%). Clinical parameters such as peripheral oxygen saturation $< 92\%$ (35.7 vs. 52.0) and those used for qSOFA calculation were more frequently abnormal at admission.

Almost all in-hospital complications were more frequent in patients older than 75 years (Figure 1). However, admission to ICU was limited to 42 patients (11.2 vs. 4.8%), and invasive mechanical ventilation was used in 4.4% of patients ≥ 75 years. Other commonly used treatments for COVID-

19, such as hydroxychloroquine, antiviral drugs, interferon and tocilizumab were prescribed less frequently to those aged 75 and over.

Analysis of the primary outcome

All-cause mortality was 541 of 1520 participants (35.6%). A higher mortality across increasing age subgroups was found, from 11.7% in patients 65-69 years to 54.5% in patients above 80 years (Figure 2). When divided into two subgroups, 65-74 and ≥ 75 years, mortality was significantly higher in the latter (18.4 vs. 48.2%, $p < 0.001$) (Appendix 1i). The influence of comorbidities in mortality in both groups is shown in Appendix 1j. A clear trend towards worse outcomes with higher CCI scores, especially in patients 75 years and older, was observed. Among patients who died invasive therapies had been used less frequently in those aged over 75 years compared to the younger group, which remained statistically significant even after adjusting for Charlson index (0 or 1 point), dementia and dependency (Appendix 1c).

We investigated those factors associated with higher risk of mortality. Univariate regression analyses are detailed in Appendix 1d. Age ≥ 75 (OR 3.54, 95% CI 1.76-8.38), Chronic kidney disease (3.36, 95% CI 1.00-11.33), dementia (8.06, 95% CI 1.45-44.85), peripheral oxygen saturation $< 92\%$ (5.85, 95% CI 2.89-11.84), severe lymphopenia (3.36, 95% CI 1.53-7.38) and qSOFA > 1 (8.31, 95% CI 2.29-30.16) were independent predictors of death (Appendix 1e and Table 3).

DISCUSSION

This is, to the best of our knowledge, one of the largest studies with detailed data focused on older patients with COVID-19 to date. As in previous studies, the main finding was a high risk of in-hospital mortality among these patients.

From the beginning of the epidemic older age has been associated with worse outcomes. For example, *Wu et al*, in the Chinese Center for Disease Control and Prevention report of 72314 cases, found an overall fatality rate of 2.4%, increasing to 8.0% in patients aged 70 to 79 years and 14.8% in those older than 80. [6] In Italy, an analysis of the first 2003 deaths from COVID-19 by the Italian National Institute of Health found that the median age of those who died was 80.5 years (IQR 74.3-85.9) with 87.7% being older than 70 years. [11] In a more recent study that included 8910 patients from Asia, Europe and America, an age greater than 65 years was found to be independently associated with an increased risk of in-hospital death (mortality of 10.0%, vs. 4.9% among those ≤ 65 years of age). [12] In our cohort of older people with COVID-19, we found an increasing in-hospital mortality across ages, from 11% in patients aged 65-69 to more than 50% in those aged 80 or older. Although this may seem an significantly higher rate of mortality than the previously mentioned studies, asymptomatic patients and those with mild degrees of the disease were not usually hospitalised and thus, they are not represented in our population of study. However, when comparing with reports of hospitalised patients from China, we also find differences in outcomes. For example, from a cohort of 1099 patients in the first two months of the outbreak, *Guan et al* reported 153 patients aged ≥ 65 with a presence of composite end-point (admission

in ICU, mechanical ventilation or death) in 20.9% of them, and death occurring in only 15 patients of the overall cohort. This may be explained by younger age of patients included (median age 47 years, with only 15.1% of patients older than 65 years). [4] A younger age in their overall population or a lower rate of older patients hospitalised in their hospitals may be behind these low percentages of older people. Supporting our belief of a higher mortality than previously reported, *McMichael et al* have recently published a rate of 33% of deaths in infected residents of a long-term care facility in Washington, US. [13]

The presence of a higher number of comorbidities and a worse immune response towards the infection may be related to older patients' susceptibility. [14] As in previous studies, we found a high prevalence of comorbidities within these patients admitted to hospital for COVID-19. Previous conditions such as cardiovascular diseases, diabetes mellitus, hypertension, chronic lung diseases, obesity, cancer and chronic kidney disease have been associated with severe illness and mortality. [5,6,12,15-17]. In our study, chronic kidney disease (eGFR<30ml/min/1.73m²) was found to be an independent risk factor of mortality, with an OR 3.36. Kidney disease has been already described as a risk factor for mortality in COVID-19. [18] Of note, SARS-Cov-2 may have itself direct cytopathic effects on kidney tissue, [3] entering kidney cells through an angiotensin converting enzyme (ACE)-2 dependent pathway. In addition, immune-mediated damage related to deposition of virus-induced immune complexes has been reported. [19] These facts may worsen function of previously damaged kidneys.

Importantly, this is one of the few studies to associate dementia with worse outcomes (OR 8.06) [20]. Globally, more than 50 million people have dementia [21] and several organizations have recently released recommendations on these patients during the outbreak. [22] We postulate that this higher risk of in-hospital death may be explained by several factors: firstly, a worse baseline health status; secondly, delayed access to hospitals, sometimes in more advanced stages of the disease due to their inability to express some initial symptoms; finally, the difficulties faced by hospital facilities in maintaining quality of care for this very dependent group of patients where isolation, PPE and health system overload are present. Furthermore, it has been recently hypothesized that the presence of ApoE e4e4 allele increases risks of severe COVID-19 infection, independent of preexisting dementia, cardiovascular disease, and type-2 diabetes. [20]. It is noteworthy that some prevention strategies have been specifically developed for patients with dementia in dedicated institutions. [23]

A very low level of lymphocytes was another independent predictor. It is known that the lymphocyte count decreases in COVID-19 patients. [24,25] Potential mechanisms may be the lymphocytes death or dysfunction caused directly by the virus or indirectly through inflammatory cytokines or metabolic molecules. [26] In our study, other important predictors of mortality were a qSOFA scale more than 1 point and a peripheral oxygen saturation less than 92% at admission, parameters which reflect a more severe degree of the disease on arrival to the emergency room.

Interestingly, patients 75 years and older were less frequently admitted to ICUs and received less therapies commonly used for COVID-19 treatment.

Lian et al, in one of the largest studies evaluating older patients' characteristics to date, reported higher rates of ICU admission (9.56% vs 1.38%) and methylprednisolone application (28.68% vs 9.36%) when compared with younger patients (cutoff point 60 years). [27] However, they did not make any distinction among ages above 60 years. Our finding of a trend towards less invasive therapies in the oldest subgroup led us to undertake a further subgroup analysis (Appendix 1c), comparing deceased patients younger and older than 75 years without significant comorbidities (Charlson Comorbidity Index of 0 or 1, without any degree of dementia or dependency). We found that there is still a subset of "previously healthy patients" who were deceased without having been admitted to any ICU or without having undergone more advanced techniques, such as mechanical ventilation, circulatory support or the use of drugs such as Tocilizumab. For this reason, we assume that age may have been the only limitation for them. This is consistent with the Italian guidelines that recommended, in resource-limited circumstances, consideration of prognosis — how long the patient is likely to live if treated —, potentially assigning a higher priority for intensive care access to younger patients with severe illness than to older patients. [28,29] However, several societies have stated that advanced age should not by itself be a criterion for excluding patients from some treatments or specialized hospital units. [13,31,32] For example, NICE guidelines recommend the use of Clinical Frailty Scale (CFS) in patients aged 65 years and older as a guidance for the decision [32], although it may also have its important limitations. [33] Furthermore, some authors consider that, for the healthcare systems in many countries, it may be crucial to reconfigure services to cope with demand. [34]

Other interesting findings from our study are the widespread use of unevicenced treatments such us hydroxychloroquine, azithromycin and antivirals; and the finding of different patterns of inital symptoms in older patients, with low levels of some of the usually considered "core features" such as anosmia. This fact should be taken into account when evaluating older patients with suspicion of COVID-19.

To conclude, long-term care facilities deserve special attention in this COVID-19 pandemic, due to their vulnerability to respiratory disease outbreaks. [35,36] It is known that they are the origin of a very high number of new cases, and a high mortality has been observed. [14] Given that vaccines and antiviral medications are not currently available for Covid-19, other prevention strategies should be enhanced. Among them, screening and restricted access for visitors, screening of health care personnel -including measurement of body temperature-, social distancing - including restricting resident movement-, staff hands-on training and education on infection prevention and PPE use, should be applied.

[13,14,37,38]

Limitations

There are some limitations to acknowledge regarding our study. Firstly, it is limited by its observational design: disparities in treatment and criteria of admission to ICU may be found among countries, centres and pandemic timing. Besides, some previous conditions and symptoms such as dementia or altered taste may be underreported, as data depended on electronic medical records. Secondly, the sample size is limited, since it only includes hospitalised patients. Thirdly, the original design of the study did not

include data about frailty, as well as presentation with delirium nor other parameters that may have been used in prognostic tools such as NEWS-2 or CURB-65. Finally, although this cohort was collected in a prospective manner, the results reported in this study are based on a post-hoc analysis and should be regarded as hypothesis-generating. However, our study provides a 'real-world' observation on the management and prognosis of these patients and interesting conclusions can be derived.

CONCLUSION

In our study, patients aged 65 years or older hospitalised for COVID-19 had high rates of in-hospital complications and mortality, especially among patients older than 75. Age ≥ 75 years, dementia, peripheral oxygen saturation at admission $< 92\%$, severe lymphopenia and qSOFA scale > 1 were independent predictors of mortality in this population. Our data supports other studies' findings that older people are more vulnerable to COVID-19, and preventive measures should be enhanced among them.

Acknowledgements: Cardiovascular Excellence SL, for their essential support in the database and registry webpage. All HOPE COVID-19 researchers.

Declaration of Conflict of interest: None.

Declaration of sources of funding: Non conditioned grant (FUNDACIÓN INTERHOSPITALARIA PARA LA INVESTIGACIÓN CARDIOVASCULAR, FIC. Madrid, Spain). This nonprofit institution had no role in the study design; in the collection, analysis, interpretation of data; in the writing of the report; nor in the decision to submit the paper for publication.

REFERENCES

Please see appendix 4 for a full list of references.

1. World Health Organization. WHO announces COVID-19 outbreak a pandemic 11-03-2020. 2020. Available in: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/events-as-they-happen> [Accessed 01-05-2020].
2. Lithander FE, Neumann S, Tenison E, Lloyd K, Welsh TJ, Rodrigues JC, et al. COVID-19 in Older People: A Rapid Clinical Review *Age and Ageing* 2020; 1–15. doi: 10.1093/ageing/afaa093
3. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395(10223):497-506. doi: 10.1016/S0140-6736(20)30183-5.
4. Guan W, Ni Z-y, Hu Y, Liang W, Ou C, He J, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. *New England Journal of Medicine*. *N Engl J Med*. 2020;382:1708-1720. doi: 10.1056/NEJMoa2002032.
5. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395(10229):1054-62. doi: 10.1016/S0140-6736(20)30566-3.
6. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72314 cases from the Chinese Center for Disease Control and Prevention. *JAMA*. 2020;323(13):1239-1242. doi: 10.1001/jama.2020.2648.

7. Shahid Z, Kalayanamitra R, McClafferty B, Kepko D, Ramgobin D, Patel R, et al. COVID-19 and Older Adults: What We Know. *J Am Geriatr Soc*. 2020;68(5):926-929. doi: 10.1111/jgs.16472.
8. International COVID19 Clinical Evaluation Registry, (HOPE COVID 19) (ClinicalTrials.gov Identifier: NCT04334291) 2020 [updated 6 April 2020; accessed 01-05-2020]. Available from: <https://clinicaltrials.gov/ct2/show/NCT04334291>.
11. Livingston E, Bucher K. Coronavirus Disease 2019 (COVID-19) in Italy. *JAMA*. 2020. doi: 10.1001/jama.2020.4344.
12. Mehra MR, Desai SS, Kuy SR, Henry TD, Patel AN. Cardiovascular Disease, Drug Therapy, and Mortality in Covid-19. *NEMJ*. 2020. DOI: 10.1056/NEJMoa2007621.
13. McMichael TM, Currie DW, Clark S, Pogosjans S, Kay M, Schwartz NG, et al. Epidemiology of Covid-19 in a Long-Term Care Facility in King County, Washington. *N Engl J Med*. 2020. doi: 10.1056/NEJMoa2005412.
14. Bonanad C, García-Blas S, Tarazona-Santabalbina FJ, Díez-Villanueva P, Ayesta A, Forés JS, Vidán-Austiz M, Formiga F, Ariza-Solé A, Martínez-Sellés M. Coronavirus: the geriatric emergency of 2020. Joint document of the Geriatric Cardiology Section of the Spanish Society of Cardiology and the Spanish Society of Geriatrics and Gerontology. *Rev Esp Cardiol*. 2020. doi: 10.1016/j.recesp.2020.03.027.
15. Liang W, Guan W, Chen R, Wang W, Li J, Xu K, et al. Cancer patients in SARS-CoV-2 infection: a nationwide analysis in China. *Lancet Oncol*. 2020;21(3):335.

16. CDC COVID-19 Response Team. Preliminary Estimates of the Prevalence of Selected Underlying Health Conditions Among Patients with Coronavirus Disease 2019—United States, February 12–March 28, 2020. *MMWR Morb Mortal Wkly Rep.* 2020. Doi: 10.1016/S1470-2045(20)30096-6.
17. Lighter J, Phillips M, Hochman S, Sterling S, Johnson D, Francois F, et al. Obesity in patients younger than 60 years is a risk factor for Covid-19 hospital admission. *Clin Infect Dis.* 2020. Doi: 10.1093/cid/ciaa415.
18. Cheng Y, Luo R, Wang K, Zhang M, Wang Z, Dong L, et al. Kidney disease is associated with in-hospital death of patients with COVID-19. *Kidney Int.* 2020. doi: 10.1016/j.kint.2020.03.005.
19. Yao XH, Li TY, He ZC, Ping YF, Liu HW, Yu SC, et al. A pathological report of three COVID-19 cases by minimally invasive autopsies. *Zhonghua Bing Li Xue Za Zhi.* 2020;49(0):9. doi: 10.3760/cma.j.cn112151-20200312-00193.
21. Alzheimer's Disease International. World Alzheimer's report 2019: attitudes to dementia. September, 2019. Available in: <https://www.alz.co.uk/research/WorldAlzheimerReport2019.pdf> (accessed April 15, 2020).
22. Chinese Society of Geriatric Psychiatry, Alzheimer's Disease Chinese, Psychogeriatric Interest Group of Chinese Society of Psychiatry, et al. Expert recommendations on mental health and psychosocial support for persons with cognitive disorders and their caregivers during the COVID-19 outbreak. *Chinese J Psychiatry* 2020; 53: 89–94.
23. Huang HT, Chen TC, Liu TY, Chiu CF, Hsieh WC, Yang CJ. How to prevent outbreak of a hospital affiliated dementia day-care facility in the

pandemic COVID-19 infection in Taiwan. *J Microbiol Immunol Infect.* 2020.

doi: 10.1016/j.jmii.2020.04.007

26. Tan L, Wang Q, Zhang D et al. Lymphopenia predicts disease severity of COVID-19: a descriptive and predictive study. *Signal Transduction and Targeted Therapy.* 2020;5. doi: 10.1038/s41392-020-0148-4.

27. Lian J, Jin X, Hao S, Cai H, Zhang S, Zheng L, et al. Analysis of Epidemiological and Clinical features in older patients with Corona Virus Disease 2019 (COVID-19) out of Wuhan. *Clin Infect Dis.* 2020. doi: 10.1093/cid/ciaa242.

28. Vergano M, Bertolini G, Giannini A, et al. Clinical Ethics Recommendations for the Allocation of Intensive Care Treatments, in Exceptional, Resource-Limited Circumstances. Italian Society of Anesthesia, Analgesia, Resuscitation, and Intensive Care (SIAARTI). 2020. Available in: [http://www.siaarti.it/SiteAssets/News/COVID19%20 - %20documenti%20SIAARTI/SIAARTI%20-%20Covid-19%20-%20Clinical%20Ethics%20Reccomendations.pdf](http://www.siaarti.it/SiteAssets/News/COVID19%20-%20documenti%20SIAARTI/SIAARTI%20-%20Covid-19%20-%20Clinical%20Ethics%20Reccomendations.pdf).

28. Emanuel EJ, Persad G, Upshur R, et al. Fair Allocation of Scarce Medical Resources in the Time of Covid-19. *N Engl J Med.* 2020. doi: 10.1056/NEJMs2005114.

29. STATEMENT of the EuGMS Executive Board on the COVID-19 epidemic. EuGMS. Available in: <https://www.eugms.org/news/read/article/489.html>. (Accessed May 01, 2020).

32. National Institute for Health and Care Excellence. COVID-19 rapid guideline: critical care. 2020. Available in:

www.nice.org.uk/guidance/ng159 (Accessed May 01, 2020).

33. Hubbard RE, Maier AB, Hilmer SN, Naganathan V, Etherton-Beer C, Rockwood K. Frailty in the Face of COVID-19. *Age and Ageing* 2020;00:1–2. doi: 10.1093/ageing/afaa095.

34. Gemelli Against COVID-19 Geriatrics Team, Landi F, Barillaro C, Bellieni A, Brandi V, Carfi A, Cipriani C, et al. The geriatrician: the frontline specialist in the treatment of COVID-19 patients. *JAMDA*. 2020. doi: 10.1016/j.jamda.2019.08.026.

37. Centers for Disease Control and Prevention. Interim infection prevention and control recommendations for patients with suspected or confirmed coronavirus disease 2019 (COVID-19) in healthcare settings. 2020. Available in: <https://www.cdc.gov/coronavirus/2019-ncov/infection-control/control-recommendations.html> (accessed 01-05-2020).

38. Centers for Disease Control and Prevention. Preparing for COVID-19: long-term care facilities, nursing homes. 2020. Available in: <https://www.cdc.gov/coronavirus/2019-ncov/healthcare-facilities/prevent-spread-in-long-term-care-facilities.html> (accessed 01-05-2020).

TABLES

Table 1. Baseline characteristics of the overall cohort

Characteristic	Overall cohort of patients ≥ 65 years (N=1520)
Age (years), median [IQR] ^a	76 [71-83]
Male sex, n (%)	916/1520 (60.3)
Race, white or caucasian race, n (%)	1402/1520 (92.2)
Body mass index (kg/m ²), median [IQR] ^b	27.6 [25.1-30.9]
Duration of stay (days), median [IQR]	8 [5-13]
Chronic conditions, n (%)	
Hypertension	1047/1514 (69.2)
Dyslipidemia	728/1499 (48.6)
Diabetes Mellitus	377/1435 (26.3)
Obesity	267/1058 (25.2)
Current Smoking	72/1520 (4.7)
Chronic kidney disease ^c	164/1440 (11.4)
Chronic lung disease	380/1502 (25.3)
Any heart disease	562/1463 (38.4)
Atrial Fibrillation	247/1518 (16.3)
Cerebrovascular disease	185/1475 (12.5)
Connective Tissue disease	63/1471 (4.3)
Liver disease	61/1456 (4.2)
Cancer	295/1464 (20.2)
Parkinsons disease	18/1154 (1.6)
Dementia	70/1154 (6.1)
Immunosuppression, n (%) ^d	119/1325 (9.0)
Prior tuberculosis, n (%)	18/1154 (1.6)
Human Immunodeficiency virus, n (%)	2/1154 (0.2)
Charlson Comorbidity Index (points), median	1 [0-3] /1011

[IQR]	
Home Oxygen Therapy, n (%)	75/1504 (5.0)
Preadmission medication, n (%) ^e	
ASA	336/1487 (22.6)
Other Antiplatelet drug	91/1456 (6.3)
Oral Anticoagulation	287/1487 (19.3)
ACEI / ARB's	743/1499 (49.6)
Beta Blockers	376/1493 (25.2)
Symptoms, n (%)	
Asymptomatic	67/1490 (4.5)
Dyspnea ^f	874/1493 (58.5)
Tachypnoea > 22 breaths per minute	450/1430 (31.5)
Fatigue	706/1448 (48.8)
Anosmia / Hyposmia	50/1376 (3.6)
Altered taste	74/1378 (5.4)
Sorethroat	136/1397 (9.7)
Fever	1165/1503 (77.5)
Cough	983/1486 (66.2)
Vomiting	109/1441 (7.6)
Diarrhea	253/1440 (17.6)
Arthromyalgia	397/1451 (27.6)
qSOFA >1, n (%)	122/1136 (10.7)
X-Ray abnormalities, n (%) ^g	1206/1520 (79.4)
Complications during admission, n (%)	
Respiratory failure	902/1501 (60.1)
Heart Failure	143/1476 (9.7)
Acute kidney Injury	369/1489 (24.8)
Upper Respiratory-Tract Infection	203/1427 (14.2)
Pneumonia	1346/1502 (89.6)
Sepsis	248/1482 (16.7)
Systemic Inflammatory Response Syndrome	342/1454 (23.5)

Any relevant bleeding ^h	38/1454 (2.6)
Hemoptysis	36/1467 (2.5)
Embolic event	19/1462 (1.3)
Oxygen Therapy at the admission, n (%)	
Oxygen during admission	1204/1506 (79.9)
High Flow Nasal Cannula	348/1480 (23.5)
Non-Invasive Mechanical Ventilation	238/1472 (15.7)
Invasive Mechanical Ventilation	92/1448 (6.4)
Prone Position	164/1454 (11.3)
Circulatory support or ECMO	41/1447 (2.8)
Duration of Mechanical Ventilation (days), median (IQR)	8.0 [3.5-13.5]
Other therapies during the Admission, n (%)	
Use of Glucocorticoids	403/1476 (27.3)
Use of chloroquine and derivates	1226/1492 (82.2)
Use of Antiviral Drugs ⁱ	899/1488 (59.1)
Use of Interferon	215/1465 (14.7)
Use of Tocilizumab	101/1466 (6.9)
Use of Antibiotics	1141/1449 (78.7)
ACEI/ARB's	329/1397 (23.6)
ICU Admission, n (%)	114/1520 (7.5)
Death, n (%)	541/1520 (35.6)

Abbreviations: **ACEI** = Angiotensin converting enzyme (ACE)

inhibitors; **ARB** = Angiotensin receptors blockers; **ASA** =

Acetylsalicylic acid; **ECMO** = Extracorporeal membrane oxygenation;

ICU = Intensive Care Unit; **qSOFA** = Quick Sequential Organ Failure Assessment Score.

^a Detailed distribution of patients according to age is shown in

Appendix 1g. ^b The body-mass index is the weight in kilograms

divided by the square of the height in meters. ^c Creatinine clearance

< 30 ml/min. ^d Immunosuppressive therapy for psoriasis arthritis, lung transplantation, kidney transplantation or systemic lupus erythematosus; oncological disease such as breast cancer, prostate cancer, myelodysplastic syndrome or gammopathy; glucocorticoid therapy caused by COPD; dialysis; HIV or hepatitis C. ^e Other premedications are listed in appendix 1b. ^f Severity of dyspnea is described in Appendix 1b. ^g Detailed data are listed in in Appendix 1b^h Rectorrhagia, hematuria, epistaxis, and popliteal aneurysm bleeding with relevant decreased hemoglobin > 2 mg/l. ⁱ Lopinavir or /and Ritonavir were the most commonly used antivirals.

UNCORRECTED MANUSCRIPT

Table 2. Characteristics of patients aged 65-74 as compared to patients aged ≥ 75 .*

Characteristic	65-74 N= 642	≥ 75 N= 878	P value
Age (years), mean \pm SD	69.9 \pm 2.8	83.2 \pm 6.1	<0.001
Male sex, n (%)	387/642 (60.3)	529/878 (60.3)	0.99
Duration of stay (days), mean \pm SD	9.4 \pm 7.3	8.6 \pm 6.9	0.05
Chronic conditions, n (%)			
Hypertension	398/639 (62.3)	649/875 (74.2)	<0.001
Dyslipidemia	294/634 (46.4)	434/865 (50.2)	0.15
Diabetes Mellitus	150/609 (24.6)	227/826 (27.5)	0.26
Obesity	132/447 (29.5)	135/611 (22.1)	<0.001
Chronic kidney disease	47/616 (7.6)	117/824 (14.2)	<0.001
Chronic lung disease	150/637 (23.5)	230/865 (26.6)	0.18
Any heart disease	156/632 (24.7)	406/858 (47.3)	<0.001
Cerebrovascular disease	46/627 (7.3)	139/848 (16.4)	<0.001
Parkinsons disease	2/513 (0.4)	16/641 (2.5)	0.01
Dementia	10/513 (1.9)	60/641 (9.4)	<0.001
Charlson Comorbidity Index	1.4 \pm 1.7	2.0 \pm 1.9	<0.001
Any dependency level	36/633 (5.7)	272/861	<0.001

		(31.6)	
Home Oxygen Therapy, n (%)	20/638 (3.1)	55/866 (6.4)	0.01
Preadmission medication, n (%)			
ASA	123/632 (19.5)	214/856 (25)	0.01
Other antiplatelet drugs	22/623 (3.5)	69/834 (8.3)	<0.001
Oral Anticoagulation	68/631 (10.8)	219/857 (25.6)	<0.001
ACEI	283/635 (44.6)	461/865 (53.3)	0.001
Beta Blockers	121/634 (19.1)	256/860 (29.8)	0.001
Beta Agonist Inhalation Therapy	74/636 (11.6)	138/849 (16.3)	0.01
Glucocorticoids Inhalation Therapy	62/634 (9.8)	118/854 (13.8)	0.02
Symptoms, n (%)			
Dyspnea	374/635 (58.9)	500/858 (58.3)	0.89
Fatigue	303/623 (48.6)	403/825 (48.8)	0.93
Fever	523/638 (82.0)	642/865 (74.2)	<0.001
Cough	437/632 (69.1)	546/854 (63.9)	0.04
Diarrhea	136/610 (22.3)	117/830 (14.1)	<0.001
Arthromyalgia	202/622	195/830	<0.001

	(32.5)	(23.5)	
Clinical parameters, n (%)			
Peripheral Oxygen Saturation < 92 %	223/625 (35.7)	445/856 (52.0)	<0.001
qSOFA Score >1, n (%)	29/500 (5.8)	93/636 (14.6)	<0.001
Laboratory parameters, n (%)			
Elevated D-Dimer	388/559 (69.4)	535/712 (75.1)	0.02
Elevated Troponin I	48/276 (17.4)	102/413 (24.7)	0.02
Severe Lymphopenia (<500/ml)	84/600 (14.0)	175/827 (21.2)	0.001
X-Ray Abnormalities	526/574 (91.6)	680/806 (84.4)	<0.001
Complications during admission			
Respiratory failure	329/636 (51.7)	573/865 (66.2)	<0.001
Heart Failure	33/626 (5.3)	110/850 (12.9)	<0.001
Acute kidney Injury	118/634 (18.6)	251/855 (29.4)	<0.001
Pneumonia	591/641 (92.2)	755/861 (87.7)	0.01
Sepsis	82/632 (13)	166/850 (19.5)	0.001
SIRS	134/628 (21.3)	208/826 (25.2)	0.09
Oxygen Therapy, n (%)			

Oxygen, any	480/635 (75.6)	724/871 (83.1)	<0.001
High Flow Nasal Cannula	121/625 (19.4)	227/855 (26.5)	0.001
Non-Invasive Mechanical Ventilation	102/623 (16.4)	136/849 (16.0)	0.86
Invasive Mechanical Ventilation	56/621 (9)	36/827 (4.4)	<0.001
Prone Position	83/615 (13.5)	81/839 (9.7)	0.02
Circulatory support or ECMO	27/620 (4.4)	14/827 (1.7)	0.01
Other therapies during the Admission, n (%)			
Use of Glucocorticoids	166/627 (26.5)	237/849 (27.9)	0.54
Use of chloroquine and derivatives	559/629 (88.9)	667/863 (77.3)	<0.001
Use of Antiviral Drugs	437/627 (69.7)	462/861 (53.7)	<0.001
Use of Interferon	105/621 (16.9)	110/844 (13.0)	0.04
Use of Tocilizumab	60/622 (9.6)	41/844 (4.9)	<0.001
Use of Antibiotics	479/610 (78.5)	662/839 (78.9)	0.86
ICU Admission, n (%)	72/642 (11.2)	42/878 (4.8)	<0.001
Death, n (%)	118/642 (18.4)	423/878 (48.2)	<0.001

Abbreviations: **ACEI** = Angiotensin converting enzyme (ACE) inhibitors; **ARB** = Angiotensin receptors blockers; **ASA** = Acetylsalicylic acid; **ECMO** = Extracorporeal membrane oxygenation;

ICU = Intensive Care Unit; **qSOFA** = Quick Sequential Organ Failure Assessment Score; **SIRS** = Systemic Inflammatory Response Syndrome.

* Extensive data are listed in Appendix 3.

UNCORRECTED MANUSCRIPT

Table 3. Summary of predictors for all cause in-hospital mortality *

	Multivariable analysis		
	OR	95%CI	P-value
Age ≥ 75	3.54	1.76-8.38	<0.001
Chronic kidney disease	3.36	1.00-11.33	0.05
Dementia	8.06	1.45-44.85	0.017
Peripheral Oxygen Saturation < 92 %	5.85	2.89-11.84	<0.001
Severe lymphopenia (<500/mm³)	3.36	1.53-7.38	0.003
qSOFA >1	8.31	2.29-30.16	0.001

qSOFA = Quick Sequential Organ Failure Assessment Score.


* A complete description of variables included in the model is shown in Appendix 1e.

FIGURES

Figure 1. Percentage of in-hospital complications in patients aged 65-74 and ≥ 75 years.

X axis = Type of complication; Y axis = Percentage (%).


* $p < 0.05$



1PT

UNCORRECTED

Figure 2. Number of patients aged ≥ 65 years hospitalised for COVID-19 (blue boxes) and percentage of deaths (red dots) according to age.



UNCORRECTED M